

Award Number: W81XWH-07-1-0012

TITLE: Cellular Mechanisms for Motor Stimulation and Neuroprotection by A2A Antagonists in PD model

PRINCIPAL INVESTIGATOR: Jiang-Fan Chen, M.D., Ph.D.

CONTRACTING ORGANIZATION: Boston University School of Medicine
Boston, MA 02118

REPORT DATE: May 2008

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE 31-05-2008		2. REPORT TYPE Annual		3. DATES COVERED 1 FEB 2007 - 31 JAN 2008	
4. TITLE AND SUBTITLE Cellular Mechanisms for Motor Stimulation and Neuroprotection by A2A Antagonists in PD model				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-07-1-0012	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Jiang-Fan Chen, M.D., Ph.D. Email: chenjf@bu.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Boston University School of Medicine Boston, MA 02118				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT None listed.					
15. SUBJECT TERMS None listed.					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			USAMRMC
U	U	U	UU	7	19b. TELEPHONE NUMBER (include area code)

BLANK PAGE

Scientific Progress Report:

Cellular Mechanisms for Motor Stimulation and Neuroprotection by A_{2A} Antagonists in PD model

1.. **Specific Aims** – not significantly modified.

2. Main findings and Significances:


Specific Aim 1: Hypothesis: A_{2A}R antagonists modulate motor activity and neuroprotection by acting at forebrain neurons and non-forebrain neurons, respectively.

Specific Aim 2: Hypothesis: A_{2A}Rs act at striatal, cerebral cortical and substantial nigral neurons to modulate MPTP-induced dopaminergic neurotoxicity.

Specific Aim 3: Hypothesis: A_{2A}R antagonists act at peripheral and brain inflammatory cells to exert neuroprotection against MPTP neurotoxicity.

C2. Forebrain- and neuron-specific deletion of the A_{2A}R gene in forebrain-A_{2A}R KO mice.

Figure C2. Forebrain- and neuron-specific inactivation of A_{2A}Rs in fb-A_{2A}R KO mice




To generate forebrain-specific A_{2A}R KO (fb-KO) mice, we cross-bred floxed A_{2A}R mice to Cre-CaMKIIα transgenic mice in which Cre proteins are expressed under the control of an 8.5 kb promoter fragment of the CaMKIIα gene^{1,2,3}. Using PCR primers targeted to the floxed A_{2A}R genes³, we detected CaMKIIα-Cre mediated A_{2A}R gene deletion (“KO” band) in Cre(+) mice, specifically in forebrain regions including striatum (ST), cerebral cortex (CT), hippocampus (HIP) and olfactory bulb (OB), but not in cerebellum (CB), brainstem (BS), or midbrain (MB) where dopaminergic neurons reside, or in six peripheral organs tested (Figure C2-A). A residual floxed band was still present in the forebrain, likely due to the A_{2A}R in glial cells that do not express CaMKIIα-Cre.

Both [³H]-SCH58261-binding (Figure C2-B) and western blot analyses (Figure C2-C) demonstrated the abolishment of A_{2A}R proteins in fb-KO to the background level observed in global-A_{2A}R KO mice (gb-KO). To further characterize the cellular specificity of the A_{2A}R gene deletion, we separated neurons (β-tubulin III⁺ cells), microglia (CD11b⁺ cells), and astrocytes (GFAP⁺ cells) from three fb-KO and two fb-WT mice (all of them with the floxed A_{2A}R) by FACS. The “KO” band was detected in the neuronal preparations from all three fb-KO mice but was not observed in the microglial or astroglial preparations from the same fb-KO mice (Figure C1-D). A residual floxed band was still present in the neuronal preparations, which may suggest the presence of striatal neurons that do not express CaMKIIα-Cre (e.g. striatal cholinergic interneurons). Additionally, [³H]-SCH58261-binding confirmed that A_{2A}R protein on microglial cells is not affected in these mice (Figure C2-E). Together, these data clearly demonstrate at the gene and protein levels that A_{2A}Rs are completely depleted in forebrain neurons in the fb-KO line.


C4. Genetic deletion of A_{2A}Rs in forebrain neurons abolishes the psychomotor effect of A_{2A}R antagonists, but does not confer neuroprotection against MPTP neurotoxicity

We evaluated the effect of the genetic deletion of A_{2A}Rs in forebrain neurons on the A_{2A}R modulation of psychomotor activity. We assessed the motor stimulant effect of the selective A_{2A}R antagonist KW-6002 and of the non-selective antagonist caffeine in fb-A_{2A}R KO mice. In accordance with previous studies³, KW-6002 (3 mg/kg) produced a strong motor stimulation in WT mice, which was completely abolished in fb-A_{2A}R-KO mice (Figure C4A). Similarly, caffeine (10 mg/kg) produced a strong motor stimulation in WT mice, which was largely attenuated in fb-A_{2A}R KO mice (Figure C4B), as occurs in gb-A_{2A}R KO mice⁴. Thus, A_{2A}Rs in forebrain neurons are critical for the A_{2A}R-mediated motor effects. We also examined the effect of global or forebrain A_{2A}Rs inactivation on MPTP-induced dopaminergic toxicity. The residual striatal dopamine levels were indistinguishable between fb-A_{2A}R KO mice and their WT littermates after a single injection of MPTP (40 mg/kg x1, i.p.) (Figure C4C) or three injections of MPTP (20 mg/kg x 2 at 2 hr interval, i.p.) (Figure 4D). In contrast, residual dopamine content in the striatum of gb-A_{2A}R KO mice was significantly higher than that of WT littermates after MPTP treatment (Figure C4D, *P<0.05), consistent with our previous study⁵. Thus, deletion of A_{2A}Rs in forebrain neurons does not confer neuroprotection against MPTP-induced brain dopaminergic toxicity.




C7. Intraventricular administration of the A_{2A}R antagonist KW-6002 into forebrain A_{2A}R-KO mice protects against MPTP neurotoxicity.

To determine if A_{2A}Rs located in brain cells other than forebrain neurons contribute to neuroprotection, we tested the effect of intracerebroventricular (icv) injection of KW-6002 on MPTP-induced dopaminergic toxicity in fb-KO mice. We reasoned that if A_{2A}Rs in forebrain neurons were the only targets of KW-6002, then KW-6002 would not be able to provide neuroprotection in fb-KO mice. If instead, KW-6002 afforded neuroprotection, it would act on A_{2A}Rs located in brain cells other than forebrain neurons. Vehicle or KW-6002 (10 µg/2 µl/brain) was administered into the left ventricle of fb-KO mice or WT littermates 10 min prior to a single injection of MPTP (40 mg/kg, i.p.). We measured dopamine content 7 days after MPTP intoxication and found that MPTP produced about a 50% reduction of striatal dopamine content in fb-KO mice and WT littermates pre-treated with vehicle (Figure C7-A). Pre-treatment with KW-6002 protected against MPTP-induced toxicity in fb-KO mice to an extent similar to that observed in WT littermates (Figure C7-A). KW-6002 also attenuated MPTP-induced reduction of TH⁺ cells in the substantia nigra in fb-KO and WT mice (Figures C7-B & C7-C). Thus, protection against MPTP-induced neurotoxicity afforded by KW-6002 is likely due to the blockade of A_{2A}Rs in brain cells other than forebrain neurons.




C8. KW-6002 markedly reduced MPTP-induced microglial activation in striatum.

To seek further evidence for the role of A_{2A} Rs in microglial cells in PD models, we investigated the influence of KW-6002 on microglial activation in the MPTP model. To better characterize microglial function *in vivo*, we have developed flow cytometry analysis to quantify striatal CD11b+ cells in different proliferation and activation stages in MPTP-intoxicated mice.



MPTP treatment. MPTP treatment increased the total CD11b+ cell numbers by 2-fold (per 5×10^5 total events) compared to saline treatment (Figures C8-A & C8-B). KW-6002 attenuated the MPTP-induced increase in CD11b+ cells. Interestingly, the most significant MPTP-induced increase in CD11b+ cells was seen in the subpopulation with a large cell size (Figure C8-B). In addition, it is in this subpopulation of large diameter CD11b+ cells that the MPTP effect is significantly attenuated by KW-6002. Thus, A_{2A} R antagonists may attenuate the over-activation rather than the initial proliferation of microglial cells. These results suggest that KW-6002 may confer neuroprotection against MPTP by attenuating microglial responses *in vivo*.

C9. Up-regulation of A_{2A} R protein in microglial cells 48 hours after MPTP treatment.



To investigate the identity of the brain cells other than forebrain neurons by which KW-6002 affords neuroprotection, we examined A_{2A} R immunoreactivity (IR) in microglial cells by double immunolabeling in MPTP-intoxicated fb-KO mice 48 hours after MPTP treatment (Figure C9). CD11b+ activated microglial cells demonstrating the characteristic cellular morphology were detected in the striatum (Figure C9-lower left panel). Double-labeling revealed that MPTP markedly induced A_{2A} R-IR in microglia (Figure C9-lower panels). As expected, no A_{2A} R-IR was detected in β -tubulin-III+ neurons in the striatum of fb-KO mice after acute MPTP treatment (Figure C9-upper middle panel). Thus, MPTP treatment markedly induces A_{2A} R expression in microglial cells in the striatum.

3. Plans:

Specific Aim 2:

- (1) Despite of the lack of neuroprotection in forebrain A_{2A} R KO mice, we are evaluating the effect of selective inactivation of the A_{2A} R in striatal neurons on MPTP-induced MPTP neurotoxicity since striatal A_{2A} Rs and cortical A_{2A} Rs have been to exert opposite effects on excitotoxicity and psychomotor behavior.
- (2) We are also performing intra-nigral injection of A_{2A} R antagonists to localize the anatomical locus whereby A_{2A} R antagonists exert neuroprotective effects.

Specific Aim 3:

- (1) We are performing bone marrow transplantation experiment to create chimeric mice with selective inactivation and reconstitution of the A2AR activity in bone marrow derived cells. Using these chimeric mice, we will critically evaluate the effect of selective manipulation of A2ARs in bone marrow-derived cells on MPTP-induced dopaminergic neurotoxicity.
- (2) We are developing microglia-specific A2A receptor knockout mice to critically evaluate the role of A2ARs in microglial cells on MPTP-induced dopaminergic neurotoxicity.

4. Publications:

[Yu L, Shen HY, Coelho JE, Araújo IM, Huang QY, Day YJ, Rebola N, Canas PM, Rapp EK, Ferrara J, Taylor D, Müller CE, Linden J, Cunha RA, **Chen JF**.](#) (2008) Adenosine A2A receptor antagonists exert motor and neuroprotective effects by distinct cellular mechanisms. Annals of Neurology. 63(3):338-46 (see attached)